

Open Research Online

The Open University's repository of research publications and other research outputs

Adhesion attenuation and enhancement in aqueous solutions

Conference or Workshop Item

How to cite:

Bowen, James; Rossetto, Herbert L. and Kendall, Kevin (2014). Adhesion attenuation and enhancement in aqueous solutions. In: Cell Adhesion Century: Culture Breakthrough, 28-29 Apr 2014, London, UK.

For guidance on citations see [FAQs](#).

© [\[not recorded\]](#)

Version: Version of Record

Link(s) to article on publisher's website:

https://www.academia.edu/9823179/Adhesion_attenuation_and_enhancement_in_aqueous_solutions

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Adhesion Attenuation and Enhancement in Aqueous Solutions

James Bowen, Hebert L Rossetto, Kevin Kendall FRS


School of Chemical Engineering, University of Birmingham, UK

Email: j.bowen.1@bham.ac.uk

1. Introduction

When two surfaces confine water layers between them at the nanoscale, the behaviour of these confined water molecules can deviate significantly from the behaviour of bulk water, which affects the adhesion of such surfaces. This study assesses the role of confined water layers on the adhesion of hydrophilic surfaces, and how sensitive this adhesion is to the presence of contaminant solutes which can disrupt the hydrogen bonding network of water molecules between the surfaces.

2. Experimental



substance	M_w (g/mol)	dielectric constant	surface tension (mN/m)	viscosity (mPa s)	refractive index
water (H ₂ O)	18.01	80.0	73.0	1.00	1.33
ethanol (≥99.5%, 200 proof, C ₂ H ₆ O)	46.07	24.3	22.1	1.20	1.36
dimethylformamide (C ₃ H ₇ NO)	73.09	38.3	25.0	0.92	1.43
formamide (CH ₃ NO)	45.04	84.0	55.2	3.30	1.44

- Alumina-sputtered sphere-tipped cantilever, interacted versus an alumina single crystal, using atomic force microscopy.
- Solvent/solute concentrations tested over the range 0-100% organic.
- Measurements were performed under immersed conditions using (i) water, and mixtures of (ii) water/dimethylformamide, (iii) water/ethanol, and (iv) water/formamide.


3. Results

$$W_{th} = -AR/6D \quad - (1)$$

$$A = \frac{3}{4}kT\left(\frac{\epsilon_1 - \epsilon_3}{\epsilon_1 + \epsilon_3}\right)^2 + \frac{3h\nu_e}{16\sqrt{2}}\frac{(n_1^2 - n_3^2)^2}{(n_1^2 + n_3^2)^{3/2}} \quad - (2)$$

A = Hamaker constant, R = tip radius, D = separation distance, k = Boltzmann's constant, T = temperature, ϵ = dielectric constant, h = Planck's constant, ν_e = electron absorption frequency, n = refractive index

- The measured work of adhesion upon separation of the surfaces was greatly in excess of the theoretical value, W_{th} , predicted by Eq. 1, assuming D = 1 nm.
- Low concentrations of dimethylformamide and ethanol increased the work of adhesion to values greater than the work of adhesion measured in water.
- Formamide and dimethylformamide exhibited differences in behaviour at high concentrations and for pure liquid.



- Viscous effects were not important for the liquids and mixtures investigated here.

4. Conclusions

- Number of possible hydrogen bonds which can be formed between surfaces strongly affects adhesion behaviour.
- Ability of atoms within the solute to act as hydrogen bond donors and acceptors is also important.
- For example, methyl hydrogen is a better donor than formyl hydrogen; the strength of a hydrogen bond depends on the specific binding environments involved.
- The number of water molecules in the primary solvation shell is important at low solute concentrations.
- Continuum theories describing adhesion between surfaces are not applicable when hydrogen bonding effects occur.
- When the two surfaces are at closest approach, the structure of the hydrogen bonded network requires further experimental and theoretical consideration.

Acknowledgements

The atomic force microscope used in this research was obtained, through Birmingham Science City: Innovative Uses for Advanced Materials in the Modern World (West Midlands Centre for Advanced Materials Project 2), with support from Advantage West Midlands (AWM) and partly funded by the European Regional Development Fund (ERDF).

This poster is a summary of the paper published in Langmuir:

"Adhesion of alumina surfaces through confined water layers containing various molecules"

Langmuir **2012**, 28, 4648-4653; DOI: 10.1021/la2045064

The URL can be obtained by scanning the adjacent QR code.

